

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 9/51, 31/192, 9/10	A1	(11) International Publication Number: WO 00/09096 (43) International Publication Date: 24 February 2000 (24.02.00)
(21) International Application Number: PCT/US99/18270 (22) International Filing Date: 12 August 1999 (12.08.99) (30) Priority Data: 09/154,422 13 August 1998 (13.08.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US US 09/154,422 (CIP) Filed on 13 August 1998 (13.08.98) (71) Applicant (for all designated States except US): NANOSYS-TEMS [US/US]; 3000 Horizon Drive, King of Prussia, PA 19406 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): LEE, Robert [US/US]; 1335 Roberts Road, Gilbertsville, PA 19525 (US). DE CASTRO, Lan [US/US]; 44 Plymouth Street, Montclair, NJ 07042 (US). (74) Agents: SCHAFER, Michele, M. et al.; Foley & Lardner, Suite 500, 3000 K Street, N.W., Washington, DC 20007-5109 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: INJECTABLE FORMULATIONS OF NANOPARTICULATE NAPROXEN (57) Abstract Described are an injectable formulation of nanoparticulate naproxen that produces minimal or no pain or burning sensation upon administration, and methods of making and using such a formulation. The injectable formulation comprises nanoparticulate naproxen having a povidone polymer adsorbed on the surface thereof in an amount sufficient to maintain an effective average particle size of less than about 600 nm.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INJECTABLE FORMULATIONS OF NANOPARTICULATE NAPROXEN

BACKGROUND OF THE INVENTION

5

Field of the Invention

The present invention is directed to an injectable formulation of nanoparticulate naproxen that produces minimal or no intramuscular pain or burning sensation upon administration, methods of making such a formulation, and methods of using such a formulation.


10

Description of the Related Art

Naproxen, also known as 6-methoxy- α -methyl-2-naphthalene-acetic acid and *d*-2-(6-methoxy-2-naphthyl)propionic acid, is a well-known anti-inflammatory, analgesic, and antipyretic agent. It has been approved in many countries around the world for almost two decades and has a very safe risk-benefit profile. It is sold under the trade names ALEVE[®], ANAPROX[®], NAPROSYN[®], and SYNFLEX[®] (all available from Syntex Chemicals, Inc.). See *The Merck Index*, 10th Edition, pp. 6274 (Merck & Co., Rahway, NJ, 1983).

20

Naproxen, which is highly water insoluble, *i.e.*, less than 10 mg/ml, has the following chemical structure:



Naproxen is a non-steroidal anti-inflammatory drug (NSAID) often used to relieve the inflammation, swelling, stiffness, and joint pain associated with rheumatoid arthritis, osteoarthritis (the most common form of arthritis), juvenile arthritis, ankylosing spondylitis (spinal arthritis), tendinitis, bursitis, and acute gout.

5 In addition, it is used to treat pain associated with menstrual periods, migraine headaches, and other types of mild to moderate pain.

Naproxen acts by suppressing the production of prostaglandins, which are hormone-like substances that act on local tissues to produce pain and inflammation. Its pharmaceutical forms of delivery include tablets, capsules, and liquids. Delivery
10 characteristics and forms are disclosed in, for example, U.S. Patent Nos. 3,904,682; 4,009,197; 4,780,320; 4,888,178; 4,919,939; 4,940,588; 4,952,402; 5,200,193; 5,354,556; 5,462,747; and 5,480,650, all of which are specifically incorporated by reference. The synthesis of naproxen is described in U.S. Patent Nos. 3,904,682 and 4,009,197.

15 Naproxen is a more potent pain reliever than aspirin, especially for menstrual cramps, toothaches, minor arthritis, and injuries accompanied by inflammation, such as tendinitis. The naproxen sodium salt is specifically indicated in the treatment of various types of acute and very high intensity pain because it induces a rapid and sustained remission. In addition, it is possible to obtain a good analgesic effect with
20 few administrations, due to naproxen's particular pharmacokinetics. Tablet formulations of naproxen were approved for OTC ("over the counter" as compared to prescription) marketing by the U.S. Food and Drug Administration in 1994.

Because of naproxen's low solubility, it is generally formulated for oral administration. However, oral administration of naproxen frequently results in
25 gastrointestinal irritation. All NSAIDs produce gastrointestinal symptoms to some degree upon oral administration. Such symptoms most commonly are constipation, gastric burns, diarrhea, stomatitis, dyspepsia, nausea, vomiting, upper abdominal

pain, and heartburn. Oral administration may also lead to an ulcer or bleeding from the stomach or duodenum.

Gastrointestinal irritation resulting from oral administration of an NSAID can be significant. Numerous literature articles detail the severity of gastric irritation
5 caused by NSAID compositions. For example, one report states that between 10,000 and 20,000 people in Canada each year are hospitalized with major gastro-intestinal bleeding caused by oral ingestion of NSAIDs, with effects resulting in death for at least 1,000 of these patients. See *Marketplace*, October 24, 1996. Yet another reference states that gastrointestinal complications of NSAID use may be responsible
10 for over 10,000 deaths each year. See *American Family Physician*, March 1997.

Injectable formulations of naproxen are preferable over oral administration forms for several reasons. First, such formulations can lessen or eliminate side effects of gastro-intestinal irritation. Second, intravenous (IV) or intramuscular (IM) administration of a drug results in a significantly shorter response time as compared
15 to oral administration. Moreover, injectable formulations of pain medication are also preferable for post-operative health care, where oral administration may not be feasible. Injectable formulations of naproxen are particularly preferred, as naproxen is not addictive, in contrast to other injectable formulations of drugs, such as morphine and ketorolac (Toradol®).

20 Injectable formulations of naproxen have been used prior to the present invention. See Marsala et al., "Treatment of Acute Pain of Ureteral and Biliary Colic with Naproxen Sodium Administered by the Parenteral Route," *Int. J. Clin. Pharmacol. Res.*, 6:495-500 (1986) (IM and IV injections of naproxen); L. Kvarnes, "Naproxen Sodium Versus Pentazocine in Treating Postoperative Pain," *Curr. Ther. Res., Clin. Exp.*, 46:259-268 (1989) (IM injections of naproxen). However,
25 injectable naproxen formulations are difficult to formulate due to the low solubility of naproxen. Moreover, current injectable formulations of naproxen are undesirable because they produce intense pain and/or a burning sensation upon administration.

5 There is currently a need for a safe and effective injectable formulation of naproxen that produces minimal or no pain or burning sensation upon administration. In addition, there is a need in the art for methods of making and methods of using such naproxen formulations. The present invention satisfies these needs.

SUMMARY OF THE INVENTION

The invention provides for compositions comprising high naproxen concentrations in low injection volumes, with rapid drug dissolution upon administration.

4

than about 600 nm. Preferably, the pH of the liquid dispersion medium is maintained within the range of from about 3 to about 8 during the size reduction process.

Yet another aspect of the present invention provides a method of treating a
5 mammal requiring anti-inflammatory, analgesic, or antipyretic treatment comprising administering to the mammal the above-described injectable nanoparticulate naproxen formulation.

It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory and are intended to
10 provide further explanation of the invention as claimed. Other objects, advantages, and novel features will be readily apparent to those skilled in the art from the following detailed description of the invention.

DETAILED DESCRIPTION OF THE INVENTION

15 The present invention is directed to the surprising and unexpected discovery of a new injectable formulation of naproxen that produces minimal or no pain or burning sensation upon administration. The injectable composition comprises nanoparticulate naproxen having a povidone polymer with a molecular weight of less than about 40,000 daltons adsorbed on the surface thereof in an amount sufficient to
20 maintain an effective average particle size of less than about 600 nm. Preferably, the effective average particle size of the nanoparticulate naproxen is less than about 450 nm, more preferably less than about 300 nm, even more preferably less than about 250 nm, and most preferably less than about 100 nm.

Naproxen has a pKa of 4.4. Below its pKa of 4.4, the solubility of naproxen
25 is about 20 micrograms/ml. However, above a pKa of 4.4, the solubility of naproxen is about 3 mg/ml. It is preferred that the nanoparticulate naproxen/povidone polymer pharmaceutical formulation of the invention has a pH of between about 6 to about 7.

In human therapy, it is important to provide a naproxen dosage form that delivers the required therapeutic amount of the drug *in vivo*, and that renders the drug bioavailable in a rapid and constant manner. The injectable nanoparticulate naproxen formulations of the present invention satisfies these needs.

5 **Povidone Polymers**

Povidone polymers, also known as polyvidon(e), povidonum, PVP, and polyvinylpyrrolidone, are sold under the trade names Kollidon® (BASF Corp.) and Plasdone® (ISP Technologies, Inc.). They are polydisperse macromolecular molecules, with a chemical name of 1-ethenyl-2-pyrrolidinone polymers and 1-vinyl-
10 2-pyrrolidinone polymers. Povidone polymers are produced commercially as a series of products having mean molecular weights ranging from about 10,000 to about 700,000 daltons. To be useful as a surface modifier for a drug compound to be administered to a mammal, the povidone polymer must have a molecular weight of less than about 40,000 daltons, as a molecular weight of greater than 40,000 daltons
15 would have difficulty clearing the body.

Povidone polymers are prepared by, for example, Reppe's process, comprising: (1) obtaining 1,4-butanediol from acetylene and formaldehyde by the Reppe butadiene synthesis; (2) dehydrogenating the 1,4-butanediol over copper at 200° to form γ -butyrolactone; and (3) reacting γ -butyrolactone with ammonia to
20 yield pyrrolidone. Subsequent treatment with acetylene gives the vinyl pyrrolidone monomer. Polymerization is carried out by heating in the presence of H₂O and NH₃. See *The Merck Index*, 10th Edition, pp. 7581 (Merck & Co., Rahway, NJ, 1983).

The manufacturing process for povidone polymers produces polymers containing molecules of unequal chain length, and thus different molecular weights.
25 The molecular weights of the molecules vary about a mean or average for each particular commercially available grade. Because it is difficult to determine the polymer's molecular weight directly, the most widely used method of classifying

various molecular weight grades is by K-values, based on viscosity measurements. The K-values of various grades of povidone polymers represent a function of the average molecular weight, and are derived from viscosity measurements and calculated according to Fikentscher's formula.

- 5 The weight-average of the molecular weight, M_w , is determined by methods that measure the weights of the individual molecules, such as by light scattering. Table 1 provides molecular weight data for several commercially available povidone polymers, all of which are soluble.

10

TABLE 1

Povidone	K-Value	Mv (Daltons)**	Mw (Daltons)**	Mn (Daltons)**
Plasdone C-15 [®]	17 ± 1	7,000	10,500	3,000
Plasdone C-30 [®]	30.5 ± 1.5	38,000	62,500*	16,500
Kollidon 12 PF [®]	11-14	3,900	2,000-3,000	1,300
Kollidon 17 PF [®]	16-18	9,300	7,000-11,000	2,500
Kollidon 25 [®]	24-32	25,700	28,000-34,000	6,000

- 15 *Because the molecular weight is greater than 40,000 daltons, this povidone polymer is not useful as a surface stabilizer for a drug compound to be administered parenterally (*i.e.*, injected).

 **Mv is the viscosity-average molecular weight, Mn is the number-average molecular weight, and Mw is the weight average molecular weight. Mw and Mn were determined by light scattering and ultra-centrifugation, and Mv was determined by viscosity measurements.

20

 Based on the data provided in Table 1, exemplary preferred commercially available povidone polymers include, but are not limited to, Plasdone C-15[®], Kollidon 12 PF[®], Kollidon 17 PF[®], and Kollidon 25[®].

Injectable Nanoparticulate Naproxen Formulations

The invention provides injectable nanoparticulate naproxen formulations that can comprise high drug concentrations in low injection volumes, with rapid drug dissolution upon administration. Preferred compositions comprise, based on % w/w:

5	naproxen	5 – 50%
	povidone polymer	0.1 – 50%
	preservatives	0.05 - 0.25%
	pH adjusting agent	pH about 6 to about 7
	water for injection	q.s.

10

Exemplary preservatives include methylparaben (about 0.18% based on % w/w), propylparaben (about 0.02% based on % w/w), phenol (about 0.5% based on % w/w), and benzyl alcohol (up to 2% v/v). An exemplary pH adjusting agent is sodium hydroxide, and an exemplary liquid carrier is sterile water for injection.

15 Other useful preservatives, pH adjusting agents, and liquid carriers are well-known in the art.

Methods of Making Injectable Naproxen Formulations

In another aspect of the invention there is provided a method of preparing the injectable nanoparticulate naproxen formulations of the invention. The method comprises: (1) dispersing naproxen in a liquid dispersion medium comprising a povidone polymer with a molecular weight of less than about 40,000 daltons; and (2) mechanically reducing the particle size of the naproxen to an effective average particle size of less than about 600 nm. Preferably, the effective average particle size of the nanoparticulate naproxen is less than about 450 nm, more preferably less than about 300 nm, even more preferably less than about 250 nm, and most preferably less than about 100 nm. The pH of the liquid dispersion medium is preferably maintained within the range of from about 5.0 to about 7.5 during the size reduction

process. Preferably, the dispersion medium used for the size reduction process is aqueous.

Effective methods of providing mechanical force for particle size reduction of naproxen include ball milling, media milling, and homogenization, for example, with a Microfluidizer[®] (Microfluidics Corp.). Ball milling is a low energy milling process that uses milling media, drug, stabilizer, and liquid. The materials are placed in a milling vessel that is rotated at optimal speed such that the media cascades and reduces the drug particle size by impaction. The media used must have a high density as the energy for the particle reduction is provided by gravity and the mass of the attrition media.

Media milling is a high energy milling process. Drug, stabilizer, and liquid are placed in a reservoir and recirculated in a chamber containing media and a rotating shaft/impeller. The rotating shaft agitates the media which subjects the drug to impaction and shear forces, thereby reducing the drug particle size.

Homogenization is a technique that does not use milling media. Drug, stabilizer, and liquid (or drug and liquid with the stabilizer added after particle size reduction) constitute a process stream propelled into a process zone, which in the Microfluidizer[®] is called the Interaction Chamber. The product to be treated is inducted into the pump, and then forced out. The priming valve of the Microfluidizer[®] purges air out of the pump. Once the pump is filled with product, the priming valve is closed and the product is forced through the interaction chamber. The geometry of the interaction chamber produces powerful forces of shear, impact, and cavitation which are responsible for particle size reduction. Specifically, inside the interaction chamber, the pressurized product is split into two streams and accelerated to extremely high velocities. The formed jets are then directed toward each other and collide in the interaction zone. The resulting product has very fine and uniform particle or droplet size. The Microfluidizer[®] also provides

a heat exchanger to allow cooling of the product. U.S. Patent No. 5,510,118, which is specifically incorporated by reference, refers to a process using a Microfluidizer[®] resulting in sub 400 nm particles.

Using a particle size reduction method, the particle size of naproxen is
5 reduced to an effective average particle size of less than about 600 nm. Preferably, the effective average particle size of the nanoparticulate naproxen is less than about 450 nm, more preferably less than about 300 nm, even more preferably less than about 250 nm, and most preferably less than about 100 nm. The naproxen particles can be reduced in size in the presence of a povidone polymer, or the povidone
10 polymer can be added to the naproxen dispersion following particle size reduction.

Naproxen can be added to a liquid medium in which it is essentially insoluble to form a premix. The concentration of the naproxen in the liquid medium can vary from about 5 to about 60%, and preferably is from about 15 to about 50% (w/v), and more preferably about 20 to about 40%. The povidone polymer can be present in the
15 premix or it can be added to the drug dispersion following particle size reduction. The concentration of the povidone polymer can vary from about 0.1 to about 50%, and preferably is from about 0.5 to about 20%, and more preferably from about 1 to about 10%, by weight.

The premix can be used directly by subjecting it to mechanical means to
20 reduce the average naproxen particle size in the dispersion to less than about 600 nm. It is preferred that the premix be used directly when a ball mill is used for attrition. Alternatively, naproxen and the povidone polymer can be dispersed in the liquid medium using suitable agitation, *e.g.*, a Cowles type mixer, until a homogeneous dispersion is observed in which there are no large agglomerates visible to the naked
25 eye. It is preferred that the premix be subjected to such a premilling dispersion step when a recirculating media mill is used for attrition.

The mechanical means applied to reduce the naproxen particle size conveniently can take the form of a dispersion mill. Suitable dispersion mills

include a ball mill, an attritor mill, a vibratory mill, and media mills such as a sand mill and a bead mill. A media mill is preferred due to the relatively shorter milling time required to provide the desired reduction in particle size. For media milling, the apparent viscosity of the premix is preferably from about 100 to about 1000
5 centipoise, and for ball milling the apparent viscosity of the premix is preferably from about 1 up to about 100 centipoise. Such ranges tend to afford an optimal balance between efficient particle size reduction and media erosion.

The attrition time can vary widely and depends primarily upon the particular mechanical means and processing conditions selected. For ball mills, processing
10 times of up to five days or longer may be required. Alternatively, processing times of less than 1 day (residence times of one minute up to several hours) are possible with the use of a high shear media mill.

The naproxen particles must be reduced in size at a temperature which does not significantly degrade naproxen. Processing temperatures of less than about 30 to
15 less than about 40°C are ordinarily preferred. If desired, the processing equipment can be cooled with conventional cooling equipment. Control of the temperature, *e.g.*, by jacketing or immersion of the milling chamber in ice water, is contemplated. Generally, the method of the invention is conveniently carried out under conditions of ambient temperature and at processing pressures which are safe and effective for
20 the milling process. Ambient processing pressures are typical of ball mills, attritor mills, and vibratory mills.

Grinding Media

The grinding media can comprise particles that are preferably substantially
25 spherical in shape, *e.g.*, beads, consisting essentially of polymeric resin. Alternatively, the grinding media can comprise a core having a coating of a polymeric resin adhered thereon.

In general, suitable polymeric resins are chemically and physically inert, substantially free of metals, solvent, and monomers, and of sufficient hardness and friability to enable them to avoid being chipped or crushed during grinding. Suitable polymeric resins include crosslinked polystyrenes, such as polystyrene crosslinked
5 with divinylbenzene; styrene copolymers; polycarbonates; polyacetals, such as Delrin[®] (E.I. du Pont de Nemours and Co.); vinyl chloride polymers and copolymers; polyurethanes; polyamides; poly(tetrafluoroethylenes), *e.g.*, Teflon[®] (E.I. du Pont de Nemours and Co.), and other fluoropolymers; high density polyethylenes; polypropylenes; cellulose ethers and esters such as cellulose acetate;
10 polyhydroxymethacrylate; polyhydroxyethyl acrylate; and silicone-containing polymers such as polysiloxanes and the like. The polymer can be biodegradable. Exemplary biodegradable polymers include poly(lactides), poly(glycolide) copolymers of lactides and glycolide, polyanhydrides, poly(hydroxyethyl methacrylate), poly(imino carbonates), poly(N-acylhydroxyproline)esters, poly(N-
15 palmitoyl hydroxyproline) esters, ethylene-vinyl acetate copolymers, poly(orthoesters), poly(caprolactones), and poly(phosphazenes). For biodegradable polymers, contamination from the media itself advantageously can metabolize *in vivo* into biologically acceptable products that can be eliminated from the body.

The grinding media preferably ranges in size from about 0.01 to about 3 mm.
20 For fine grinding, the grinding media is preferably from about 0.02 to about 2 mm, and more preferably from about 0.03 to about 1 mm in size.

The polymeric resin can have a density from about 0.8 to about 3.0 g/cm³.

In a preferred grinding process the particles are made continuously. Such a method comprises continuously introducing naproxen into a milling chamber,
25 contacting the naproxen with grinding media while in the chamber to reduce the naproxen particle size, and continuously removing the nanoparticulate naproxen from the milling chamber.

The grinding media is separated from the milled nanoparticulate naproxen using conventional separation techniques, in a secondary process such as by simple filtration, sieving through a mesh filter or screen, and the like. Other separation techniques such as centrifugation may also be employed.

5

Particle Size

As used herein, particle size is determined on the basis of the weight average particle size as measured by conventional particle size measuring techniques well known to those skilled in the art. Such techniques include, for example, sedimentation field flow fractionation, photon correlation spectroscopy, light scattering, and disk centrifugation. By "an effective average particle size of less than about 600 nm" it is meant that at least 90% of the particles, by weight, have a particle size of less than about 600 nm when measured by the above-noted techniques. In preferred embodiments, the effective average particle size is less than about 450 nm, and more preferably less than about 400 nm. The naproxen particles can also have an effective average particle size of less than about 300 nm, less than about 250 nm, and less than about 100 nm. With reference to the effective average particle size, it is preferred that at least 90%, more preferably at least 95%, and most preferably at least 99% of the particles have a particle size less than the effective average particle size. In particularly preferred embodiments essentially all of the particles have a size less than about 600 nm.


While applicants do not wish to be bound by theoretical mechanisms, it is believed that the povidone polymer hinders the flocculation and/or agglomeration of the naproxen particles by functioning as a mechanical or steric barrier between the particles, minimizing the close, interparticle approach necessary for agglomeration and flocculation.

Method of Treatment

Yet another aspect of the present invention provides a method of treating a mammal, including a human, requiring anti-inflammatory, analgesic, or antipyretic treatment comprising administering to the mammal the injectable nanoparticulate naproxen formulation of the invention. Particularly advantageous features of the present invention include that the pharmaceutical formulation of the invention exhibits unexpectedly rapid drug absorption upon administration, and produces minimal or no pain or irritation upon administration. In addition, the injectable formulation of the invention can provide a high naproxen concentration in a small volume to be injected. A general protocol for administration comprises a bolus injection of naproxen, with one continuous fast injection, rather than a slow infusion of the drug.

Sterile Product Manufacturing

Development of injectable compositions requires the production of a sterile product. The manufacturing process of the present invention is similar to typical known manufacturing processes for sterile suspensions. A typical sterile suspension manufacturing process flowchart is as follows:



As indicated by the optional steps in parentheses, some of the processing is dependent upon the method of particle size reduction and/or method of sterilization. For example, media conditioning is not required for a milling method that does not
5 use media. If terminal sterilization is not feasible due to chemical and/or physical instability, aseptic processing can be used.

The following examples are given to illustrate the present invention. It should be understood, however, that the invention is not to be limited to the specific conditions or details described in these examples

10 **Example 1**

The purpose of this example was to evaluate the use of different types of surface modifiers for the preparation of an injectable nanoparticulate formulation of naproxen.

15 The study was conducted by screening eleven surface stabilizers to identify the most suitable stabilizer for parenteral administration of naproxen. The dispersions were formulated at 40% solids to 2.4% surface stabilizer.

TABLE 2

Surface Stabilizer	Manufacturer	Results
Plasdone C15 [®] (polyvinylpyrrolidone)	ISP Technologies, Inc.	size reduction < 400 nm
Kollidon 17PF [®] (a polyvinylpyrrolidone polymer)	BASF Corp.	size reduction < 400 nm
Povidone K30 [®] (a polyvinylpyrrolidone polymer)	ISP Technologies, Inc.	size reduction < 400 nm
Tyloxapol	Nycomed, Inc.	size reduction > 400 nm
Pluronic F68 [®] (a high molecular weight polyoxyalkylene ether)	BASF Corp.	size reduction > 400 nm
Pluronic F108 [®] (a high molecular weight polyoxyalkylene ether)	BASF Corp.	size reduction > 400 nm
Tween 80 [®] (a polyoxyethylene sorbitan fatty acid ester)	ICI Americas	size reduction > 400 nm
dioctylsulfosuccinate (CAS No. 577-11-7; aka Docusate Sodium)	Ashland Chem. Co., Columbus, OH	size reduction > 400 nm
B20-5000 [®] (a triblock copolymer surface modifier)	Dow Chemical	size reduction > 400 nm
B20-5000-sulfonated (a triblock copolymer surface modifier)	Dow Chemical	size reduction > 400 nm
lecithin (CAS No. 8002-43-5)	Ashland Chem. Co., Columbus, OH	size reduction > 400 nm
Povidone K30 [®] and Pluronic F108 [®]	ISP Technologies, Inc. and BASF Corp.	size reduction > 400 nm

Only the use of the povidone polymers Plasdone C15[®], Povidone K30[®], and Kollidon 17PF[®] resulted in a nanoparticulate naproxen composition having an effective average particle size of less than about 400 nm.

Example 2

The purpose of this example was to determine the rates of absorption for intravenously and intramuscularly administered naproxen.

The plasma concentration of naproxen ($\mu\text{g/ml}$) in rabbits after administration of 15 mg/kg by either the IV or IM routes versus time in hours was determined. The method used for quantification of naproxen in rabbit plasma was modified from a procedure by Shimek et al., *Journal of Pharm. Sci.*, 15:436-439 (1982).

For IM administration, the mean peak plasma level concentration (C_{max}) of $45 \pm 9 \mu\text{g/ml}$ was reached at two hours. The absorption after IM injection was rapid with a mean plasma concentration of $23.5 \pm 5.1 \mu\text{g/ml}$ at 30 minutes after injection. At twelve hours post injection, plasma concentrations were still measurable at $5.2 \pm 3.4 \mu\text{g/ml}$.

Pharmacokinetic Analysis: The area-under-curve value (AUC) (*i.e.*, plasma-concentration time curve) calculations for the IM and IV routes were $287.1 \mu\text{g/ml} \cdot \text{hr}$ and $212.0 \mu\text{g/ml} \cdot \text{hr}$, respectively. Mean clearance was calculated to be 3.4 ml/min after IM administration and 5.1 ml/min following IV administration. The calculated $T_{1/2}$ (the time period in which the maximum plasma concentration drops by half) following IM administration was 6.9 hours, and 7.0 hours following IV administration.

20

Example 3

The purpose of this example was to demonstrate the feasibility of using a Microfluidizer[®] for aseptically producing an injectable formulation of a naproxen colloidal dispersion.

A Microfluidizer[®], Model No. M110EH (Microfluidics Corp.), was successfully used to produce a sterile colloidal dispersion of naproxen suitable for injection. The naproxen slurry was heat sterilized prior to microfluidizing.

Example 4

The purpose of this example was to determine the potential local irritation of an injectable nanoparticulate naproxen formulation when administered
5 intramuscularly to rats. This information correlates to the irritation or pain experienced upon administration of the formulation.

Test Formulation

Injectable nanoparticulate naproxen having a concentration of 489 mg/ml was used, comprising 489 mg/ml naproxen, 40 mg/ml of ISP Plasdone™ C15, 1.9 mg/ml
10 methylparaben (preservative), and 0.2 mg/ml propylparaben. Dosages of 49 and 134 mg/kg were administered intramuscularly. The intramuscular route was chosen because it is a possible route of administration to human. The test and control articles were stored at room temperature and protected from light.

Test animals

15 Twenty male Albino rats (Sprague-Dawley Crl:CD®) were used for the study, obtained from Charles River. The animals were laboratory bred and were experimentally naïve at the outset of the study. Animals selected for use were as uniform in age and weight as possible. they were approximately 7-8 weeks of age, and their body weights ranged from 210.7 to 247.0 gm. Each animal was identified
20 by a unique number via an ear tag. All animals were acclimated to laboratory conditions for approximately 12 days prior to study initiation, and the animals were housed individually in stainless-steel cages.

PMI Feeds, Inc.®, Certified Rodent Chow was available *ad libitum* via food
hoppers. No contaminants are known to be present in the certified diet that would
25 interfere with the results of the study. Tap water was available *ad libitum* via automatic watering device or water bottle. The water was routinely analyzed for contaminants. No contaminants are known to be present in the water that would interfere with the results of the study.

All animals were kept in one room and with no additional studies or other species housed in the same room. The room was well ventilated (>10 air changes per hour) with 100% fresh air (no air recirculation). A 12-hour light/12-hour dark photoperiod was maintained. Room temperature and relative humidity were set to be maintained between 22±3°C and 40 to 70%, respectively.

Administration Protocol and Methods

The twenty test animals were assigned to treatment groups as shown in Table 3 below. Each animal received a single intramuscular injection of the test article and the vehicle in distinct previously shaven sites of contralateral legs: thigh muscle of the right and left hind legs, respectively. Prior to the day of dosing (designated Day 1 of the study), all animals were fasted overnight. Following dosing, animals were returned to their cages and subsequently provided with hoppers containing food. The animals were evaluated for changes in clinical signs and body weight and the injection sites were examined for any local reaction. Ten animals each were sacrificed at approximately 48 and 96 hours after the IM injection and the injection site areas removed and processed for histopathological examination.

Table 3: Treatment Groups					
Group	Number of Males	Treatment	Dose (mg/kg)	Dose Vol. (ml/kg)	Day of Sacrifice
1	5	Control	0	0.1	3
		Nanonaproxen	49	0.1	
2	5	Control	0	0.1	5
		Nanonaproxen	49	0.1	
3	5	Control	0	0.275	3
		Nanonaproxen	134	0.275	
4	5	Control	0	0.275	5
		Nanonaproxen	134	0.275	

Antemortem Procedures

Animals were observed once daily prior to dosing. During the study, each animal was observed once each morning and afternoon for changes in general appearance and behavior. In addition, the injection sites were examined daily thereafter for local reactions. The severity of any injection site observation was graded and if possible measured (length, width, and height). Body weights were measured prior to dosing and at sacrifice.

1. Clinical Observations and Mortality

Twice daily individual clinical observations are presented in Table 4. No animals died and no treatment-related clinical signs were seen in animals given 49 mg/kg of nanoparticulate naproxen by intramuscular injection. At a dose of 134 mg/kg/ treatment-related signs such as chromorhinorrhea, pallor, rough coat, and some chromodacryorrhea and brown staining were observed.

Table 4: Clinical Observations		
Test Article	Animal No.	Clinical Observations*
Nanoparticulate Naproxen; 49 mg/kg For 3 day test period	311M	Days 1-3: normal
	312M	Days 1-3: normal
	313M	Days 1-3: normal
	314M	Days 1-2: normal Day 3: abrasion neck right side; stain red neck right side; scab scapula right
	315M	Days 1-3: normal
Nanoparticulate Naproxen; 49 mg/kg For 5 day test period	321M	Days 1-5: normal
	322M	Days 1-5: normal
	323M	Day 1: normal Day 2: slight bluish color at injection site 2 Days 3-5: normal
	324M	Days 1-4: normal Day 5: dehydrated; scab at injection site on right hind limb
	325M	Days 1-5: normal
Nanoparticulate Naproxen; 134 mg/kg	331M	Day 1: normal Days 2-3: infrequent stool Day 3: chromorhinorrhea, pallor

Table 4: Clinical Observations		
Test Article	Animal No.	Clinical Observations*
For 3 day test period	332M	Day 1: normal Days 2-3: infrequent stool; chromorhinorrhea; pallor; rough coat Day 3: chromodacryorrhea-both; brown-stained scrotum
	333M	Day 1: normal Days 2-3: infrequent stool Day 3: chromorhinorrhea; pallor; dehydrated; scab at right injection site
	334M	Day 1: normal Days 2-3: infrequent stool; chromorhinorrhea; wet coat pelvic region Day 3: pallor; scabs at both injection sites left and right hind limbs
Nanoparticulate Naproxen; 134 mg/kg For 5 day test period	335M	Day 1: normal Days 2-3: infrequent stool Day 3: chromodacryorrhea both; chromorhinorrhea; dehydrated; pallor; diarrhea; dark stool
	341M	Day 1: normal Days 2-4: chromorhinorrhea Days 2-5: infrequent stool; pallor Day 3: brown-stained forefeet Days 3-5: rough coat Day 4: prostrate Day 5: dehydrated; scab at injection sites on right hand limb
	342	Day 1: normal Day 2: chromorhinorrhea Days 2-3: infrequent stool Days 3-5: rough coat; pallor Days 4-5: scab head Day 5: scab at injection sites on left hand limb
	343M	Day 1: normal Days 2-5: infrequent stool Days 3-5: piloerection; rough coat; dehydrated; pallor
	344M	Day 1: normal Day 2: loose stool; chromorhinorrhea Days 2-3: infrequent stool Days 2-4: rough coat Day 5: scabs at injection sites bilaterally
	345M	Day 1: normal Days 2-5: infrequent stool; chromorhinorrhea; pallor Day 3: rough coat; chromodacryorrhea both Days 3-5: brown-stained forefeet Day 5: dehydrated; scabs at injection sites on hind limbs

*Findings exclude mechanical artifacts (clipper abrasions).

2. Body Weight

Individual and group mean body weights are presented in Table 5. The Day 3 low-dose group (49 mg/kg) was inadvertently not fasted for necropsy and, therefore, showed a much larger weight gain than the fasted low-dose group sacrificed on Day 5. However, animals in the high-dose group (134 mg/kg) sacrificed on Days 3 (also, inadvertently not fasted) or 5 showed an overall average weight loss that was considered to be treatment related.

Table 5: Individual and Group Mean Body Weights (g)				
Treatment	Animal No.	Day 1	Day 3*	Day 5
Nanoparticulate Naproxen; 49 mg/kg	311M	210.7	251.5	
	312M	219.4	256.6	
	313M	224.5	260.9	
	314M	234.7	271.8	
	315M	237.3	273.0	
	mean	225.3	262.8	
	±SD	±11.0	±9.4	
Nanoparticulate Naproxen; 49 mg/kg	321M	218.9		233.5
	322M	216.8		222.4
	323M	223.4		229.5
	324M	234.6		249.1
	325M	241.9		259.3
	mean	227.1		238.8
	±SD	±10.8		±15.1
Nanoparticulate Naproxen; 134 mg/kg	331M	215.2	203.2	
	332M	216.6	210.1	
	333M	226.7	211.0	
	334M	228.0	216.3	
	335M	247.0	227.3	
	mean	226.7	213.6	
	±SD	±12.7	±9.0	
Nanoparticulate Naproxen; 134 mg/kg	341M	216.1		192.1
	342M	213.0		184.5
	343M	226.3		189.7
	344M	228.3		208.5
	345M	241.5		209.2
	mean	225.0		196.8
	±SD	±11.3		±11.3

*Terminal body weight from unfasted animals.

3. Injection Site Evaluation

Individual and group mean dermal irritation (erythema/eschar) and (edema) are presented in Tables 6 and 7, respectively. No erythema/eschar or edema responses were noted in any of the animals (indicated by "0" in Tables 6 and 7).

Table 6: Individual and Group Mean Dermal Irritation (Erythema or Eschar) Scores					
Treatment	Animal No.	Day 2	Day 3	Day 4	Day 5
Control	311M	0	0		
	312M	0	0		
	313M	0	0		
	314M	0	0		
	315M	0	0		
	mean	0	0		
	±SD	±0	±0		
Nanoparticulate Naproxen; 49 mg/kg	311M	0	0		
	312M	0	0		
	313M	0	0		
	314M	0	0		
	315M	0	0		
	mean	0	0		
	±SD	±0	±0		
control	321M	0	0	0	0
	322M	0	0	0	0
	323M	0	0	0	0
	324M	0	0	0	0
	325M	0	0	0	0
	mean	0	0	0	0
	±SD	±0	±0	±0	±0
Nanoparticulate Naproxen; 49 mg/kg	321M	0	0	0	0
	322M	0	0	0	0
	323M	0	0	0	0
	324M	0	0	0	0
	325M	0	0	0	0
	mean	0	0	0	0
	±SD	±0	±0	±0	±0
Control	331M	0	0		
	332M	0	0		
	333M	0	0		
	334M	0	0		
	335M	0	0		

Table 6: Individual and Group Mean Dermal Irritation (Erythema or Eschar) Scores					
Treatment	Animal No.	Day 2	Day 3	Day 4	Day 5
	mean	0	0		
	±SD	±0	±0		
Nanoparticulate Naproxen; 134 mg/kg	331M	0	0		
	332M	0	0		
	333M	0	0		
	334M	0	0		
	335M	0	0		
	mean	0	0		
	±SD	±0	±0		
control	341M	0	0	0	0
	342M	0	0	0	0
	343M	0	0	0	0
	344M	0	0	0	0
	345M	0	0	0	0
	mean	0	0	0	0
	±SD	±0	±0	±0	±0
Nanoparticulate Naproxen; 134 mg/kg	341M	0	0	0	0
	342M	0	0	0	0
	343M	0	0	0	0
	344M	0	0	0	0
	345M	0	0	0	0
	mean	0	0	0	0
	±SD	±0	±0	±0	±0

Table 7: Individual and Group Mean Dermal Irritation (Edema) Scores					
Treatment	Animal No.	Day 2	Day 3	Day 4	Day 5
Control	311M	0	0		
	312M	0	0		
	313M	0	0		
	314M	0	0		
	315M	0	0		
	mean	0	0		
	±SD	±0	±0		
Nanoparticulate Naproxen; 49 mg/kg	311M	0	0		
	312M	0	0		
	313M	0	0		
	314M	0	0		
	315M	0	0		
	mean	0	0		
	±SD	±0	±0		

Table 7: Individual and Group Mean Dermal Irritation (Edema) Scores					
Treatment	Animal No.	Day 2	Day 3	Day 4	Day 5
Control	321M	0	0	0	0
	322M	0	0	0	0
	323M	0	0	0	0
	324M	0	0	0	0
	325M	0	0	0	0
	mean	0	0	0	0
	±SD	±0	±0	±0	±0
Nanoparticulate Naproxen; 49 mg/kg	321M	0	0	0	0
	322M	0	0	0	0
	323M	0	0	0	0
	324M	0	0	0	0
	325M	0	0	0	0
	mean	0	0	0	0
	±SD	±0	±0	±0	±0
control	331M	0	0		
	332M	0	0		
	333M	0	0		
	334M	0	0		
	335M	0	0		
	mean	0	0		
	±SD	±0	±0		
Nanoparticulate Naproxen; 134 mg/kg	331M	0	0		
	332M	0	0		
	333M	0	0		
	334M	0	0		
	335M	0	0		
	mean	0	0		
	±SD	±0	±0		
control	341M	0	0	0	0
	342M	0	0	0	0
	343M	0	0	0	0
	344M	0	0	0	0
	345M	0	0	0	0
	mean	0	0	0	0
	±SD	±0	±0	±0	±0
Nanoparticulate Naproxen; 134 mg/kg	341M	0	0	0	0
	342M	0	0	0	0
	343M	0	0	0	0
	344M	0	0	0	0
	345M	0	0	0	0
	mean	0	0	0	0
	±SD	±0	±0	±0	±0

Postmortem Procedures

All animals were terminated by CO₂ asphyxiation and exsanguination. Gross necropsy was limited to examination of the injection sites. No treatment-related effects were observed as all injection sites (muscle) were normal. One animal (#333) had a single red subcutaneous focus above the muscle injection site that probably corresponded to hemorrhage. Injection site scabs noted on the twice daily clinical observations generally corresponded to dried blood. Although no internal gross examination was performed, two high-dose animals (#'s 332 and 333) were observed to have severely distended stomachs filled with fluid and gas when they were terminated by exsanguination. These observed findings were considered to be treatment related.

Conclusion

No dermal irritation was observed following single-dose intramuscular administration of injectable nanoparticulate naproxen at doses of 49 mg/kg and 134 mg/kg, in comparison to the control animals. Similarly, no treatment-related clinical signs were observed at doses of 49 mg/kg. At a dose of 134 mg/kg, the following clinical signs were observed: chromorhinorrhea, pallor, rough coat, some chromodacryorrhea, and brown staining.

At necropsy, no treatment-related gross findings at the injection sites were observed. Similarly, histopathological examination of the injection sites revealed no treatment-related effects.

* * * * *

It will be apparent to those skilled in the art that various modifications and variations can be made in the compositions, methods, and uses of the present invention without departing from the spirit or scope of the invention. Thus, it is intended that the present invention cover the modifications and variations of this

invention provided they come within the scope of the appended claims and their equivalents.

WE CLAIM:

1. An injectable pharmaceutical composition comprising: (1) naproxen particles having an effective average particle size of less than about 600 nm; (2) a
5 povidone polymer adsorbed on the surface of the naproxen particles; and (3) a pharmaceutically acceptable carrier.
2. The composition of claim 1, wherein the effective average particle size of the naproxen particles is less than about 450 nm.
10
3. The composition of claim 1, wherein the effective average particle size of the naproxen particles is selected from the group consisting of less than about 400 nm, less than about 300 nm, less than about 250 nm, and less than about 100 nm.
4. The composition of claim 1, wherein the povidone polymer is present
15 in an amount of about 0.1 to about 50% (w/w) based on the total weight of the naproxen and povidone polymer.
5. The pharmaceutical composition of claim 1, wherein the povidone
20 polymer has a molecular weight of about 40,000 daltons.
6. The pharmaceutical composition of claim 1, wherein the povidone polymer has a molecular weight of less than about 40,000 daltons.
7. The composition of claim 1, wherein the naproxen is present in an
25 amount of about 5.0 to about 50% (w/w), by weight.

8. A method of treating a mammal comprising administering to the mammal an effective amount of an injectable pharmaceutical composition comprising: (1) naproxen particles having an effective average particle size of less than about 600 nm; (2) a povidone polymer adsorbed on the surface of the naproxen particles; and (3) a pharmaceutically acceptable carrier.

9. The method of claim 8, wherein the effective average particle size of the naproxen particles is less than about 450 nm.

10. The method of claim 9, wherein the effective average particle size of the naproxen particles is selected from the group consisting of less than about 400 nm, less than about 300 nm, less than about 250 nm, and less than about 100 nm.

11. The method of claim 8, wherein the povidone polymer has a molecular weight of about 40,000 daltons.

12. The method of claim 8, wherein the povidone polymer has a molecular weight of less than about 40,000 daltons.

13. A method of preparing an injectable pharmaceutical composition that produces minimal or no pain or irritation upon administration, wherein the composition comprises: (1) naproxen particles having an effective average particle size of less than about 600 nm; (2) a povidone polymer adsorbed on the surface of the naproxen particles; and (3) a pharmaceutically acceptable carrier, wherein the method comprises:

(a) dispersing the naproxen particles in a liquid dispersion medium comprising a povidone polymer; and

- (b) mechanically reducing the particle size of the naproxen to an effective average particle size of less than about 600 nm.

14. The method of claim 13, wherein the effective average particle size of the naproxen particles is less than about 450 nm.

15. The method of claim 14, wherein the effective average particle size of the naproxen particles is selected from the group consisting of less than about 400 nm, less than about 300 nm, less than about 250 nm, and less than about 100 nm.

16. The method of claim 13, wherein the povidone polymer has a molecular weight of about 40,000 daltons.

17. The method of claim 13, wherein the povidone polymer has a molecular weight of less than about 40,000 daltons.

18. A method of preparing an injectable pharmaceutical composition comprising: (1) naproxen particles having an effective average particle size of less than about 600 nm; (2) a povidone polymer adsorbed on the surface of the naproxen particles; and (3) a pharmaceutically acceptable carrier, wherein the method comprises:

- (a) dispersing the naproxen particles in a liquid dispersion;
- (b) mechanically reducing the particle size of the naproxen to an effective average particle size of less than about 600 nm; and
- (c) adding a povidone polymer to the dispersion of naproxen particles.

19. The method of claim 18, wherein the effective average particle size of the naproxen particles is less than about 450 nm.

20. The method of claim 19, wherein the effective average particle size of the naproxen particles is selected from the group consisting of less than about 400 nm, less than about 300 nm, less than about 250 nm, and less than about 100 nm.

5

21. The method of claim 18, wherein the povidone polymer has a molecular weight of about 40,000 daltons.

22. The method of claim 18, wherein the povidone polymer has a
10 molecular weight of less than about 40,000 daltons.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/18270

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K9/51 A61K31/192 A61K9/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 510 118 A (BOSCH ET AL.) 23 April 1996 (1996-04-23) column 3, line 65 -column 4, line 26 column 9; table 1	1-22
A	EP 0 577 215 A (STERLING WINTHROP INC.) 5 January 1994 (1994-01-05) page 2, line 1 - line 32 page 11; example 17	1-22
A	EP 0 463 228 A (ALFA WASSERMANN S.P.A.) 2 January 1992 (1992-01-02) claim 1	1-22
A	GB 2 046 094 A (A. NATTERMANN & CIE GMBH) 12 November 1980 (1980-11-12) page 2; example 3 claims 1-10	1-22

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

21 December 1999

Date of mailing of the international search report

12/01/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Benz, K

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/18270

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 8-12 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims: it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

information on patent family members

Inte: onal Application No

PCT/US 99/18270

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5510118	A	23-04-1996	AU 4867396 A WO 9625152 A	04-09-1996 22-08-1996
EP 577215	A	05-01-1994	US 5399363 A AU 675432 B AU 4156093 A CA 2098242 A CN 1084391 A CZ 9301316 A FI 933040 A HU 64832 A JP 7165562 A MX 9303950 A NO 932403 A NZ 248042 A SG 55089 A SK 68193 A US 5494683 A	21-03-1995 06-02-1997 06-01-1994 02-01-1994 30-03-1994 16-02-1994 02-01-1994 28-03-1994 27-06-1995 31-01-1994 03-01-1994 26-10-1994 21-12-1998 02-02-1994 27-02-1996
EP 463228	A	02-01-1992	IT 1242642 B AU 627639 B AU 6769190 A CA 2031994 A JP 4005230 A NZ 236356 A PT 96263 A US 5096926 A US 5128373 A	16-05-1994 27-08-1992 24-10-1991 18-10-1991 09-01-1992 27-01-1993 31-10-1991 17-03-1992 07-07-1992
GB 2046094	A	12-11-1980	DE 2914788 A AT 369651 B AU 539044 B AU 5733180 A BE 882731 A CA 1144480 A CH 642546 A FR 2453644 A IT 1148790 B JP 1671396 C JP 3037524 B JP 55141407 A NL 8002097 A SE 451427 B SE 8002714 A US 4309421 A ZA 8002157 A	16-10-1980 25-01-1983 06-09-1984 16-10-1980 31-07-1980 12-04-1983 30-04-1984 07-11-1980 03-12-1986 12-06-1992 05-06-1991 05-11-1980 14-10-1980 12-10-1987 12-10-1980 05-01-1982 27-05-1981